

The Group Inverse of extended Symmetric and Periodic Jacobi Matrices

S. Gago

Abstract. In this work, we explicitly compute the group inverse of symmetric and periodic Jacobi matrices with constant elements that have been extended by adding a row and a column conveniently defined. For this purpose, we interpret such matrices as the combinatorial Laplacian of a non-complete wheel that has been obtained by adding a vertex to a cycle and some edges conveniently chosen. The obtained group inverse is an incomplete block matrix with a block Toeplitz structure. In addition, we obtain the effective resistances and the Kirchhoff index of non-complete wheels.

1. INTRODUCTION AND NOTATION

The invertibility of nonsingular tridiagonal or block tridiagonal matrices has been studied in recent years; see [13] for a review on the subject. Moreover, explicit inverses are known in some cases, for instance when the tridiagonal matrix is symmetric with constant diagonals and subject to some restrictions. In [1], da Fonseca and Petronilho obtained explicit inverses of tridiagonal 2-Toeplitz and 3-Toeplitz matrices which generalize some well-known results concerning the inverse of a tridiagonal Toeplitz matrix. Recently, M.J. Jiménez in her PhD Thesis gave an explicit formula for the inverse of (p, r) -Toeplitz Jacobi matrices (tridiagonal matrices whose diagonals are formed by quasi periodic sequences, see [11],[12]). The techniques used in the mentioned results are mainly based on the theory of orthogonal polynomials. For the singular case, there is also a big amount of work, for instance in [4] the authors carried out an exhaustive analysis of the generalized inverses of singular irreducible symmetric M -matrices. The key idea of the approach was to identify any symmetric M -matrix with a positive semi-definite Schrödinger operator on a connected network whose conductances are given by the off-diagonal elements of the M -matrix. Moreover, explicit expressions for the group inverses in the cases of tridiagonal matrices and circulant matrices were obtained [6].

In this work, we present a new formula for the group inverse of an extended symmetric and periodic Jacobi matrix with constant elements. By extended we mean that a row and a column have been added to the matrix. The idea is to see the extended symmetric and periodic Jacobi matrix as the combinatorial Laplacian of a non-complete wheel and hence to obtain the group inverse, by using some previous results obtained in [7]. A non-complete wheel is a wheel where the central vertex is connected to a few vertices of the cycle. This kind of networks has many applications in Computer Science as the central vertex is called a hub (see [2]). Therefore, the non-complete wheel can be seen as a cycle with an added vertex and some new edges and hence, the result of [7] can be applied. Finally, we use the formula for the group inverse to give the effective resistances and the Kirchhoff Index a non-complete wheel.

In the following, the triple $\Gamma = (V, E, c)$ denotes a finite network, that is, a finite graph without loops nor multiple edges, with vertex set $V = \{1, \dots, n\}$ and edge set E , where each edge $e_{ij} = \{i, j\}$ has associated a conductance $c_{ij} > 0$. The standard inner product on \mathbb{R}^n is denoted by $\langle \cdot, \cdot \rangle$, thus, if $u, v \in \mathbb{R}^n$, then $\langle u, v \rangle = \sum_{k=1}^n u_k v_k$. For any $i = 1, \dots, n$ we denote by e_i the i -th vector of the standard basis of \mathbb{R}^n , by \mathbf{j} the all-ones vector of dimension n and by \mathbf{J} the all-ones matrix.

A matrix is called a *block Toeplitz matrix* iff is a block matrix, which contains blocks that are repeated in each descending diagonal from left to right, as a Toeplitz matrix in which each descending diagonal from left to right is constant. A matrix A of order $n = md + p$, $0 < p < d$, is called *incomplete block matrix* if it

is partitioned from the top-left-hand corner using $d \times d$ submatrices as far as possible. Thus

$$\mathbf{A} = \begin{bmatrix} \mathbf{A}_{11} & \cdots & \mathbf{A}_{1m} & \mathbf{A}_{1m+1} \\ \vdots & \ddots & \vdots & \vdots \\ \mathbf{A}_{m1} & \cdots & \mathbf{A}_{mm} & \mathbf{A}_{mm+1} \\ \mathbf{A}_{m+11} & \cdots & \mathbf{A}_{m+1m} & \mathbf{A}_{m+1m+1} \end{bmatrix},$$

where \mathbf{A}_{ij} , $i, j = 1, 2, \dots, m$ is an $d \times d$ matrix, \mathbf{A}_{im+1} and \mathbf{A}_{m+1j}^\top , $i, j = 1, 2, \dots, m$ are $d \times p$ matrices and \mathbf{A}_{m+1m+1} is an $p \times p$ matrix.

The combinatorial Laplacian or simply the Laplacian of the network Γ is the matrix $\mathbf{L} = (\mathbf{L}_{ij})$, where $\mathbf{L}_{ii} = \sum_{k=1}^n c_{ik}$, and $\mathbf{L}_{ij} = -c_{ij}$ when $i \neq j$. It is well-known that the Laplacian is singular, symmetric and positive semi-definite. Moreover $\mathbf{L}\mathbf{u} = 0$ iff \mathbf{u} is proportional to vector \mathbf{j} . The group inverse of the Laplacian, $\mathbf{L}^\#$, is known as the Green matrix of the network \mathbf{G} .

For every pair of vertices $\{i, j\}$ we define the dipole between them as the vector $\pi_{ij} = \mathbf{e}_i - \mathbf{e}_j$. Observe that $\pi_{ii} = 0$ and $\pi_{ij} = \pi_{ji}$. The *effective resistance* between two vertices $\{i, j\}$ of a network Γ , is defined as

$$(1) \quad R(i, j) = \langle \mathbf{G}\pi_{ij}, \pi_{ij} \rangle = (\mathbf{G})_{ii} + (\mathbf{G})_{jj} - 2(\mathbf{G})_{ij},$$

and the total resistance of the network or *Kirchhoff index* is defined as

$$K(\Gamma) = \frac{1}{2} \sum_{i,j \in V} R(i, j) = n \sum_{i=1}^n (\mathbf{G})_{ii}.$$

Given $m > 1$, $d \geq 1$, we consider a cycle C_n on $n = md$ vertices with constant conductances $c = c(i, i+1) > 0$ for any $i = 1, \dots, n-1$. A *non-complete wheel* (n, m) - W , is a network obtained from C_n by adding a new vertex $n+1$ to m of the vertices of the cycle placed at the same distance, d , with new conductances $a = c(n+1, 1+d(i-1))$, for any $i = 1, \dots, m$.

It is known that the Laplacian matrix of the cycle C_n , is a circulant matrix $\mathbf{L} = \text{circ}(2c, -c, 0, \dots, 0, -c)$ and its group inverse is

$$(\mathbf{G})_{ij} = \frac{1}{12cn} (n^2 - 1 - 6|i-j|(n-|i-j|)), \quad i, j = 1, \dots, n,$$

see [4].

2. THE GROUP INVERSE OF NON-COMPLETE WHEELS

In this section we give an explicit expression for the group inverse of extended symmetric and periodic Jacobi matrices. Not surprisingly, the expression is an incomplete block Toeplitz matrix whose coefficients involve Chebyshev polynomials. In order to obtain the claimed expression we consider the Laplacian matrix of the non-complete wheel network (n, m) - W in terms of the Laplacian of the base cycle. Then, we obtain the matrix of order $n+1$ given by

$$\mathbf{L}' = \begin{bmatrix} \mathbf{L} + \mathbf{D} & \mathbf{s} \\ \mathbf{s}^\top & \alpha \end{bmatrix},$$

where \mathbf{D} is a diagonal matrix whose non null elements, a , are placed at the $(1+d(i-1))$ -elements of the diagonal, $i = 1, \dots, m$, $\mathbf{s} = -a \sum_{i=1}^m \mathbf{e}_{1+d(i-1)}$ and $\alpha = ma$.

Theorem 1. *The group inverse of \mathbf{L}' is*

$$(\mathbf{L}')^\# = \begin{pmatrix} \mathbf{L}'_{11} & \mathbf{L}'_{12} \\ (\mathbf{L}'_{12})^\top & \mathbf{L}'_{22} \end{pmatrix},$$

where

$$\begin{aligned} \mathbf{L}'_{22} &= \frac{12cdn + an(d^2 - 1)}{12ac(n+1)^2}, \\ \mathbf{L}'_{12} &= -\frac{d}{a(n+1)^2} \left(\frac{a}{12cd}(d^2 - 1) + n + 2 \right) \mathbf{n} \otimes \mathbf{j}_m, \end{aligned}$$

with

$$(\mathbf{n})_i = -\frac{a}{12cn} \left(n(6+d) + 5m + 6mi(i-d-2) \right), \text{ for any } i = 1, \dots, d,$$

and where \mathbf{L}'_{11} is the block Toeplitz matrix

$$\mathbf{L}'_{11} = \begin{bmatrix} \mathbf{N}_1 & \mathbf{N}_2 & \dots & \mathbf{N}_m \\ \mathbf{N}_m & \mathbf{N}_1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \mathbf{N}_2 \\ \mathbf{N}_2 & \dots & \mathbf{N}_m & \mathbf{N}_1 \end{bmatrix}$$

such that any submatrix \mathbf{N}_k , $k = 1, \dots, m$, has entries

$$\begin{aligned} (\mathbf{N}_k)_{i \ h+1} &= \frac{1}{2cn} \left(-|i-1-k(d-1)-h|(n-|i-1-k(d-1)-h|) \right. \\ &\quad + \frac{1}{T_m(q)-1} \left[n(h-i+1)(V_{k-1}(q) - V_{m-k}(q)) \left(\frac{an}{c}(i-1)(h-d) - nd \right) (U_{k-2}(q) + U_{m-k}(q)) \right. \\ &\quad \left. \left. - \frac{2cd}{a}(V_{m-1}(q) - 1) - d^2 U_{m-1}(q) \right] + dh + nh - nd(2kd - 3d + 2h - n)(i-1) + kd(n-2h) \right. \\ &\quad \left. - d^2(k-1)^2 + \frac{1}{n+1} \left(\frac{(n-1)(d^2-1)}{6} - (i-2-d)i - h^2 + dh - d - 1 \right) \frac{a(d^2-1) + 12cd}{6a(n+1)^2} \right), \end{aligned}$$

with $i = 1, \dots, d$ $h = 0, \dots, d-1$ and $q = \frac{ad}{2c} + 1$.

Proof. In order to obtain $(\mathbf{L}')^\#$, we use [7, Corollary 1] which reads:

$$(\mathbf{L}')^\# = \frac{n^2}{\alpha^2(1+n)^2} \begin{pmatrix} \mathbf{C}\mathbf{M}\mathbf{C}^\top + \frac{\alpha}{n^2}\mathbf{J}_n & -\frac{\alpha}{n}\mathbf{j} - \mathbf{C}\mathbf{M}\mathbf{s} \\ -\frac{\alpha}{n}\mathbf{j}^\top - \mathbf{s}^\top\mathbf{M}\mathbf{C}^\top & \alpha + \mathbf{s}^\top\mathbf{M}\mathbf{s} \end{pmatrix},$$

where

$$\mathbf{C} = \frac{1}{n} (\alpha(n+1)\mathbf{I}_n + \mathbf{j}\mathbf{s}^\top),$$

and $\mathbf{M} = \mathbf{G} - \mathbf{G}\mathbf{\Pi}(\mathbf{I} + \mathbf{\Pi}^\top\mathbf{G}\mathbf{\Pi})^{-1}\mathbf{\Pi}^\top\mathbf{G}$, with $\mathbf{\Pi}$ the matrix whose columns are the dipoles

$$\sqrt{\frac{a}{m}} \pi_{1+(i-1)d \ 1+(j-1)d}, \quad \text{for any } 1 \leq i < j \leq m.$$

So, to obtain the result we basically need to compute matrix

$$\mathbf{M} = \mathbf{G} - \mathbf{G}\mathbf{\Pi}(\mathbf{I} + \mathbf{\Pi}^\top\mathbf{G}\mathbf{\Pi})^{-1}\mathbf{\Pi}^\top\mathbf{G}.$$

To do this we need to calculate the different involved matrices and their products in different steps. For the sake of readability, we develop the proofs in Section 4, throughout some technical lemmas.

- i) In order to obtain $B = (I + \Pi^\top G \Pi)^{-1}$, we use a reduced form

$$\Pi^\top G \Pi = \Pi_R^\top G_R \Pi_R,$$

where G_R is the $m \times m$ submatrix of $G - \frac{n^2-1}{12cn}J$ whose rows and columns are placed at $1 + (k-1)d$, for any $k = 1, \dots, m$, and Π_R is the submatrix of Π obtained only by the m rows placed at $1 + (k-1)d$, for any $k = 1, \dots, m$

- ii) Now we apply Woodbury's formula, see [15], to get

$$B = (I + \Pi_R^\top G_R \Pi_R)^{-1} = I - \Pi_R^\top (G_R^{-1} + \Pi_R \Pi_R^\top)^{-1} \Pi_R.$$

Thus, M becomes

$$\begin{aligned} M &= G - G \Pi [I - \Pi_R^\top (G_R^{-1} + \Pi_R \Pi_R^\top)^{-1} \Pi_R] \Pi^\top G \\ &= G - G [\Pi \Pi^\top - \Pi \Pi_R^\top (G_R^{-1} + \Pi_R \Pi_R^\top)^{-1} \Pi_R \Pi^\top] G = \\ &= G - G [\Pi \Pi^\top - \Pi \Pi_R^\top M_R \Pi_R \Pi^\top] G, \end{aligned}$$

where $M_R = (G_R^{-1} + \Pi_R \Pi_R^\top)^{-1}$.

Observe that both G_R and $\Pi_R^\top \Pi_R$ are, respectively, the circulant matrices

$$G_R = -\frac{d}{2cm} \text{circ}(0, m-1, \dots, j(m-j), \dots, m-1),$$

$$\Pi_R^\top \Pi_R = \frac{a}{m} (mI - J) = \frac{a}{m} \text{circ}(m-1, -1, \dots, -1).$$

Thus, for computing G_R^{-1} and $G_R^{-1} + \Pi_R^\top \Pi_R$, we use Lemma 1.

- iii) Now, from [6, Theorem 3.5], we obtain M_R in Lemma 2.
- iv) We next consider $F = \Pi \Pi^\top - \Pi \Pi_R^\top M_R \Pi_R \Pi^\top$. Notice that $\Pi \Pi^\top$ is a matrix which has the circulant matrix $\frac{a}{m} \text{circ}(m-1, -1, \dots, -1)$ as a submatrix at the rows and columns placed at $1 + (k-1)d$, for any $k = 1, \dots, m$, and 0 otherwise. It turns out that, F is the Toeplitz matrix described in Lemma 3.
- v) Since, $M = G - GFG$, we first compute matrix $H = GF$, see Lemma 4.
- vi) Next, we compute matrix $K = GFG = HG$ in Lemma 5.
- vii) Again, M is a block Toeplitz matrix whose expression is obtained in Proposition 2.
- viii) Finally, using Lemma 6, the claimed result of Theorem 1 follows.

□

3. EFFECTIVE RESISTANCES AND KIRCHHOFF INDEX FOR NON-COMPLETE WHEELS

The Green matrix is a fundamental tool for computing some desired parameters of the network, like the effective resistances of the network or the Kirchhoff index, very useful in electric circuit theory or in organic chemistry, as natural indexes describing important structural properties of circuits or molecules. Therefore, once we have compute the Green matrix, the effective resistances between two vertices of the new network are easily obtained from Formula (1) obtained in [3].

Proposition 1. *Given two vertices of Γ' , the effective resistances of the network between them are:*

- a) if $i, j \neq n+1$, $i = (k_1 - 1)d + h_1$, $j = (k_2 - 1)d + h_2$ and $k' = k_2 - k_1 + 1$, where $1 \leq k_1 \leq k_2 \leq n$ and $h_1, h_2 = 0, \dots, d-1$, with $h_1 < h_2$ when $k_1 = k_2$, then

$$\begin{aligned} R(i, j) = & \frac{-1}{T_m(q) - 1} \left[\frac{1}{c} (h_2 - h_1) (V_{k'-1}(q) - V_{m-k'}(q)) \left(\frac{a}{c^2} h_1 (h_2 - d) - \frac{d}{c} \right) (U_{k'-2}(q) + U_{m-k'}(q)) \right. \\ & + \left. \left(\frac{a}{2c^2} (h_1^2 + h_2^2 - (h_1 + h_2)d) - \frac{d}{c} \right) U_{m-1}(q) \right] + \frac{1}{cn} \left(|h_1 - h_2 - k'(d-1)| (n - |h_1 - h_2 - k'(d-1)|) \right. \\ & - (2k'd + 2h_2)h_1 - k'd(n - 2h_2) + d^2(k' - 1)^2 + h_1^2 + h_2^2 + (3h_1 - 2h_2)d + n(h_1 - h_2 + d) \Big) \\ & - \frac{(n-1)^2(d^2-1)}{6c(n+1)^2}, \end{aligned}$$

- b) if $i = (k_1 - 1)d + h_1 \neq n+1$ and $j = n+1$, then

$$\begin{aligned} R(i, n+1) = & -\frac{1}{T_m(q) - 1} \left[\frac{d}{an} (V_{m-1}(q) - 1) \frac{1}{2cn} \left(\frac{an}{c} (h_1^2 - h_1d) - nd + d^2 \right) U_{m-1}(q) \right] \\ & + \frac{h_1^2 - h_1d}{c(n+1)^2} + \frac{12cd(n^2+1) + an^2(d^2-1)}{12acn(n+1)^2} \frac{(6h_1^2 - 6h_1d - 1)}{6c(n+1)^2} \left(\frac{a(d^2-1)}{12cd} + n + 2 \right), \end{aligned}$$

with $q = \frac{ad}{2c} + 1$. Moreover, the Kirchoff index of Γ' is

$$K(\Gamma') = -\frac{1}{2c} \frac{n+1}{T_m(q) - 1} \left[\left(\frac{an}{6c} - dn + d^2 \right) U_{m-1}(q) + \left(\frac{2cd}{a} + \frac{dn}{3} \right) (V_{m-1}(q) - 1) \right] + \frac{d}{a} + \frac{dn(n+1)}{6c} - \frac{(d-1)^2n}{12c}.$$

Proof. The proof is straightforward if we observe that for $i, j \neq n+1$, then $i = (k_1 - 1)d + h_1$, $j = (k_2 - 1)d + h_2$ for any $1 \leq k_1 \leq k_2 \leq n$ and $h_1, h_2 = 0, \dots, d-1$, with $h_1 < h_2$ when $k_1 = k_2$, and if $k' = k_2 - k_1 + 1$, then

$$R(i, j) = (\mathbf{N}_1)_{h_1+1, h_1+1} + (\mathbf{N}_1)_{h_2+1, h_2+1} - 2(\mathbf{N}_{k_2-k_1+1})_{h_1+1, h_2+1},$$

and for $i = (k_1 - 1)d + h_1 \neq n+1$ and $j = n+1$, then

$$R(i, n+1) = (\mathbf{N}_1)_{h_1+1, h_1+1} + \mathbf{L}'_{22} - 2(\mathbf{L}'_{12})_{h_1+1},$$

and simplifying we obtain the claimed result. Similarly, the Kirchoff index of the non-complete wheel (n, m) - W is obtained by simplifying the expression

$$K(\Gamma') = (n+1)\text{trace}(\mathbf{L}') = (n+1)\text{trace}(\mathbf{L}'_{11}) + (n+1)\mathbf{L}'_{22}.$$

□

Finally, if we consider the case where $d = 1$, $n = m$, we obtain a complete Wheel, and in this case we notice that the Kirchhoff index coincides with the result obtained in [3].

Corollary 1. *The Kirchhoff index of a Wheel on $n+1$ vertices is*

$$K(\Gamma') = -\frac{1}{2c} \frac{n+1}{T_n(q) - 1} \left[\left(\frac{an}{6c} - n + 1 \right) U_{n-1}(q) + \left(\frac{2c}{a} + \frac{n}{3} \right) (V_{n-1}(q) - 1) \right] \frac{1}{a} + \frac{n(n+1)}{6c}.$$

4. TECHNICAL LEMMAS

In this section we include the technical lemmas and their proofs. Observe that the first Lemma excludes the case of adding a pendant vertex ($m = 1$). Besides, we point out that the result of the second Lemma 2 is obtained from the application of Theorem 3.5. from [6].

Lemma 1. *The inverse matrix of G_R is the circulant matrix*

$$G_R^{-1} = \frac{12c}{n(m^2 - 1)} \text{circ}(b_0, -b_1, -1, \dots, -1, -b_1),$$

where $b_0 = \frac{m^3 - m - 6}{6}$ and $b_1 = \frac{m^3 - m + 12}{12}$. On the other hand

$$G_R^{-1} + \Pi_R^T \Pi_R = \frac{1}{n(m^2 - 1)} \text{circ}(c_0, -c_1, -c_2, \dots, -c_2, -c_1),$$

where

$$c_0 = 12cb_0 + ad(m - 1)(m^2 - 1), c_1 = 12cb_1 + ad(m^2 - 1), c_2 = 12c + ad(m^2 - 1).$$

Lemma 2. *The circulant matrix $G_R^{-1} + \Pi_R^T \Pi_R$ is invertible iff $m > 1$ and in this case*

$$M_R = (G_R^{-1} + \Pi_R^T \Pi_R)^{-1} = \text{circ}(m_1, \dots, m_m),$$

where

$$m_j = \left[\frac{[U_{j-2}(q) + U_{m-j}(q)]d}{2c[T_m(q) - 1]} - \frac{12c + ad(m^2 - 1)}{12acm} \right], \quad j = 1, \dots, m,$$

with $q = \frac{ad}{2c} + 1$.

Lemma 3. *The matrix F is a block Toeplitz matrix given by*

$$F = \begin{bmatrix} F_1 & F_2 & \dots & F_m \\ F_m & F_1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & F_2 \\ F_2 & \dots & F_m & F_1 \end{bmatrix}$$

where each submatrix F_i has all its elements equal 0 except the first one, and

$$(F_i)_{11} = f_i = a\delta_{i1} - \frac{a^2d}{2c} \frac{[U_{i-2}(q) + U_{m-i}(q)]}{[T_m(q) - 1]}, \text{ for any } i = 1, \dots, m,$$

with $q = \frac{ad}{2c} + 1$.

Proof. Observe that

$$\begin{aligned} f_1 &= \frac{a(m-1)}{m} + \frac{a^2}{m} \sum_{j=1}^m m_j - a^2 m_1, \\ f_i &= -\frac{a}{m} + \frac{a^2}{m} \sum_{j=1}^m m_j - a^2 m_i, \text{ for any } i = 2, \dots, d. \end{aligned}$$

Therefore, we first compute

$$\begin{aligned}
\sum_{j=1}^m m_j &= \sum_{j=1}^m \left[\frac{[U_{j-2}(q) + U_{m-j}(q)]d}{2c[T_m(q) - 1]} - \frac{12c + ad(m^2 - 1)}{12acm} \right] \\
&= \frac{d}{2c[T_m(q) - 1]} \left[\sum_{j=1}^m U_{j-2}(q) + U_{m-j}(q) \right] - \frac{12c + ad(m^2 - 1)}{12ac} \\
&= \frac{d}{2c[T_m(q) - 1]} \frac{2c}{ad} [T_m(q) - 1] - \frac{12c + ad(m^2 - 1)}{12ac} = \frac{-d}{12c} (m^2 - 1).
\end{aligned}$$

And now we have

$$f_1 = \frac{a(m-1)}{m} - \frac{a^2 d(m^2 - 1)}{12cm} - a^2 \left[\frac{[U_{m-1}(q)]d}{2c[T_m(q) - 1]} - \frac{12c + ad(m^2 - 1)}{12acm} \right] = a - \frac{a^2 d}{2c} \frac{U_{m-1}(q)}{[T_m(q) - 1]},$$

and for any $i = 2, \dots, d$,

$$f_i = -\frac{a}{m} - \frac{a^2 d(m^2 - 1)}{12cm} - a^2 \left[\frac{[U_{i-2}(q) + U_{m-i}(q)]d}{2c[T_m(q) - 1]} - \frac{12c + ad(m^2 - 1)}{12acm} \right] = -\frac{a^2 d}{2c} \frac{[U_{i-2}(q) + U_{m-i}(q)]}{[T_m(q) - 1]}.$$

□

Lemma 4. *The matrix \mathbf{H} is a block Toeplitz matrix given by*

$$\mathbf{H} = \begin{bmatrix} \mathbf{H}_1 & \mathbf{H}_2 & \dots & \mathbf{H}_m \\ \mathbf{H}_m & \mathbf{H}_1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \mathbf{H}_2 \\ \mathbf{H}_2 & \dots & \mathbf{H}_m & \mathbf{H}_1 \end{bmatrix}$$

where each submatrix \mathbf{H}_k has all its elements equal 0 except its first column, and for any $i = 1, \dots, d$

$$\begin{aligned}
(\mathbf{H}_1)_{i1} &= -\frac{d}{n} - \frac{a}{2c} \frac{1}{[T_m(q) - 1]} \left[(i-1)(V_{m-1}(q) - 1) - dU_{m-1}(q) \right], \quad \text{and for any } k = 2, \dots, d, \\
(\mathbf{H}_k)_{i1} &= -\frac{d}{n} - \frac{a}{2c} \frac{1}{[T_m(q) - 1]} \left[(i-1)(V_{k-1}(q) - V_{m-k}(q)) \frac{d}{c} [a(i-1) + c] (U_{k-2}(q) + U_{m-k}(q)) \right],
\end{aligned}$$

with $q = \frac{ad}{2c} + 1$.

Proof. We point out that \mathbf{G} is also a block Toeplitz matrix and thus \mathbf{H} is again a block Toeplitz matrix, and hence we need to compute just the first d rows. We first define the following summations:

$$s_1(i, j) = \sum_{p=i}^j f_p, \quad s_2(i, j) = \sum_{p=i}^j p f_p \quad \text{and} \quad s_3(i, j) = \sum_{p=i}^j p^2 f_p.$$

In particular,

$$\begin{aligned}
K_1 &= s_1(1, m) = a - \frac{a}{2} \frac{1}{[T_m(q) - 1]} 2 [T_m(q) - 1] = 0, \\
K_2 &= s_2(1, m) = -\frac{a}{2} \frac{m}{[T_m(q) - 1]} [V_{m-1}(q) - 1], \\
K_3 &= s_3(1, m) = -\frac{2c}{d} - \frac{a}{2} \frac{1}{[T_m(q) - 1]} \left[m^2 V_{m-1}(q) - 2m U_{m-2}(q) - m(m+2) \right].
\end{aligned}$$

And moreover,

$$\begin{aligned} B_1 &= s_1(k+1, m) = -\frac{a}{2} \frac{1}{[T_m(q) - 1]} \left[V_{m-1}(q) - V_{k-1}(q) + V_{m-k}(q) - 1 \right], \\ B_2 &= s_2(k+1, m) = -\frac{a}{2} \frac{1}{[T_m(q) - 1]} \left[mV_{m-1}(q) + k(V_{m-k}(q) - V_{k-1}(q)) \right. \\ &\quad \left. + U_{k-2}(q) + U_{m-k}(q) - U_{m-2}(q) - (m+1) \right]. \end{aligned}$$

Now observe that when $j = 1 + (k-1)d$ and $k = 1, \dots, m$, for $i = 1, \dots, d$ we have that

$$\begin{aligned} (\mathbf{H}_k)_{i1} &= \sum_{\ell=1}^k g_{i,1+(\ell-1)d} f_{k-\ell+1} + \sum_{\ell=k+1}^m g_{i,1+(\ell-1)d} f_{m+k-\ell+1} g_{i,1} f_k + \sum_{\ell=2}^k g_{i,1+(\ell-1)d} f_{k-\ell+1} \\ &\quad + \sum_{\ell=k+1}^m g_{i,1+(\ell-1)d} f_{m+k-\ell+1} = A_1 + A_2 + A_3. \end{aligned}$$

Firstly we suppose $k > 1$. We compute separately A_2 and A_3 as follows

$$\begin{aligned} A_{21} &= \sum_{\ell=2}^k f_{k-\ell+1} = \sum_{p=1}^{k-1} f_p = s_1(1, k-1), \\ A_{22} &= \sum_{p=1}^{k-1} (k-p) f_p = j s_1(1, k-1) - s_2(1, k-1), \\ A_{23} &= \sum_{p=1}^{k-1} (k-p)^2 f_p = k^2 s_1(1, k-1) - 2k s_2(1, k-1) + s_3(1, k-1), \end{aligned}$$

and thus,

$$\begin{aligned} A_2 &= \sum_{\ell=2}^k g_{i,1+(\ell-1)d} f_{k-\ell+1} = \frac{1}{12cn} \left[(n^2 - 1 + 6(i-1)(n+i-1)) A_{21} - 6d(n+2i-2) A_{22} + 6d^2 A_{23} \right] \\ &= \frac{1}{12cn} \left[\alpha_1 A_{21} + \alpha_2 A_{22} + \alpha_3 A_{23} \right] = \frac{1}{12cn} \left[(\alpha_1 + \alpha_2 k + \alpha_3 k^2) s_1(1, k-1) - (\alpha_2 + 2\alpha_3 k) s_2(1, k-1) \right. \\ &\quad \left. + \alpha_3 s_3(1, k-1) \right]. \end{aligned}$$

Besides,

$$\begin{aligned} A_{31} &= \sum_{\ell=k+1}^m f_{m+k-\ell+1} = \sum_{p=k+1}^m f_p = s_1(k+1, m), \\ A_{32} &= \sum_{\ell=k+1}^m (\ell-1) f_{m+k-\ell+1} = (m+k) s_1(k+1, m) - s_2(k+1, m), \\ A_{33} &= \sum_{\ell=k+1}^m (\ell-1)^2 f_{m+k-\ell+1} = (m+k)^2 s_1(k+1, m) - 2(m+k) s_2(k+1, m) + s_3(k+1, m), \end{aligned}$$

and thus,

$$\begin{aligned}
A_3 &= \sum_{\ell=k+1}^m g_{i,1+(\ell-1)d} f_{m+k-\ell+1} = \frac{1}{12cn} [\alpha_1 A_{31} + \alpha_2 A_{32} + \alpha_3 A_{33}] \\
&= \frac{1}{12cn} [(\alpha_1 + \alpha_2(m+k) + \alpha_3(m+k)^2) s_1(k+1, m) - (\alpha_2 + 2\alpha_3(m+k)) s_2(k+1, m) + \alpha_3 s_3(k+1, m)] \\
&= \frac{1}{12cn} [(\alpha_1 + \alpha_2 k + \alpha_3 k^2) s_1(k+1, m) - (\alpha_2 + 2\alpha_3 k) s_2(k+1, m) + \alpha_3 s_3(k+1, m) \\
&\quad + (\alpha_2 m + \alpha_3 m^2 + 2\alpha_3 m k) B_1 - 2m\alpha_3 B_2].
\end{aligned}$$

Now adding all $A_1 + A_2 + A_3 =$

$$\begin{aligned}
&= \frac{1}{12cn} [(n^2 - 1 - 6(i-1)(n-i+1)) f_k + (\alpha_1 + \alpha_2 k + \alpha_3 k^2)(-f_k) - (\alpha_2 + 2\alpha_3 k)(K_2 - k f_k) \\
&\quad + \alpha_3(K_3 - k^2 f_k) - 2m\alpha_3 B_2 + (\alpha_2 m + \alpha_3 m^2 + 2\alpha_3 m k) B_1] \\
&= \frac{1}{12cn} [(n^2 - 1 - 6(i-1)(n-i+1) - \alpha_1) f_k - (\alpha_2 + 2\alpha_3 k) K_2 + \alpha_3 K_3 - 2m\alpha_3 B_2 \\
&\quad + (\alpha_2 m + \alpha_3 m^2 + 2\alpha_3 m k) B_1] \\
&= -\frac{d}{n} - \frac{a}{2c} \frac{1}{[T_m(q) - 1]} \left[(i-1)(V_{k-1}(q) - V_{m-k}(q)) - \frac{d}{c} (a(i-1) + c)(U_{k-2}(q) + U_{m-k}(q)) \right].
\end{aligned}$$

Now observe that if $k = 1$ then $j = 1$, and in this case $A_2 = 0$. Besides it holds $s_1(2, m) = -f_1$, $s_2(2, m) = K_2 - f_1$ and $s_3(2, m) = K_3 - f_1$.

Thus

$$\begin{aligned}
(H_1)_{i1} &= g_{i,1} f_1 + \sum_{\ell=2}^m g_{i,1+(\ell-1)d} f_{m+2-\ell} \\
&= \frac{1}{12cn} [(n^2 - 1 - 6(i-1)(n-i+1)) f_1 - (\alpha_1 + \alpha_2(m+1) + \\
&\quad \alpha_3(m+1)^2) f_1 - (\alpha_2 + 2\alpha_3(m+1))(K_2 - f_1) + \alpha_3(K_3 - f_1)] \\
&= \frac{1}{12cn} [(n^2 - 1 - 6(i-1)(n-i+1) - \alpha_1 - \alpha_2(m+1) - \alpha_3(m+1)^2) \\
&\quad + \alpha_2 + 2\alpha_3(m+1) - \alpha_3) f_1 - (\alpha_2 + 2\alpha_3(m+1)) K_2 + \alpha_3 K_3] \\
&= \frac{1}{12cn} [(n^2 - 1 - 6(i-1)(n-i+1) - \alpha_1 - \alpha_2 m - \alpha_3 m^2) f_1 \\
&\quad - (\alpha_2 + 2\alpha_3(m+1)) K_2 + \alpha_3 K_3] \\
&= \frac{1}{12cn} [(12d(i-1) - 6dn - 12d^2) K_2 + 6d^2 K_3] \\
&= -\frac{d}{n} - \frac{a}{2c} \frac{1}{[T_m(q) - 1]} [(i-1)(V_{m-1}(q) - 1) - dU_{m-1}(q)].
\end{aligned}$$

□

Lemma 5. *The matrix K is a block Toeplitz matrix given by*

$$K = \begin{bmatrix} K_1 & K_2 & \dots & K_m \\ K_m & K_1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & K_2 \\ K_2 & \dots & K_m & K_1 \end{bmatrix}$$

where each submatrix K_k has as elements

$$\begin{aligned} (K_k)_{i,h+1} = & -\frac{1}{2cn} \frac{1}{T_m(q) - 1} \left[n(h-i+1)(V_{k-1}(q) - V_{m-k}(q)) - \left(\frac{an}{c}(i-1)(h-d) - nd \right) (U_{k-2}(q) + U_{m-k}(q)) \right. \\ & - \frac{2cd}{a} (V_{m-1}(q) - 1) - d^2 U_{m-1}(q) \Big] + dh + nh - nd - \frac{n^2}{6} + \frac{d^2}{6} + (2kd - 3d + 2h - n)(i-1) \\ & + kd(n-2h) - d^2(k-1)^2, \end{aligned}$$

with $q = \frac{ad}{2c} + 1$, for any $k = 1, \dots, m$, $i = 1, \dots, d$ and $h = 0, \dots, d-1$.

Proof. Again we multiply two block Toeplitz matrices and thus K is a block Toeplitz matrix, and hence we can just compute only the first d rows. For each submatrix K_k , $k = 1, \dots, m$, we compute its elements. We point out that the $i(h+1)$ -element of K_k is the ij -element of K , with $j = 1 + (k-1)d + h$, for any $h = 0, \dots, d-1$.

$$\begin{aligned} (K_k)_{i,h+1} &= \sum_{\ell=1}^m h_{i,1+(\ell-1)d} \cdot g_{1+(\ell-1)d,j} = g_{i,1} h_{i,1} + \sum_{\ell=2}^k h_{i,1+(\ell-1)d} \cdot g_{1+(\ell-1)d,j} + \sum_{\ell=k+1}^m h_{i,1+(\ell-1)d} \cdot g_{1+(\ell-1)d,j} \\ &= A_1 + A_2 + A_3. \end{aligned}$$

We compute separately

$$\begin{aligned} A_1 &= \frac{1}{12cn} h_{i,1} [n^2 - 1 - 6(h+kd-d)(n-h-kd+d)], \\ A_2 &= \frac{1}{12cn} \sum_{\ell=2}^k h_{i,1+(\ell-1)d} [n^2 - 1 - 6(h+kd-\ell d)(n-h-kd+\ell d)] \\ &= \frac{1}{12cn} \sum_{\ell=2}^k h_{i,1+(\ell-1)d} [n^2 - 1 - 6(-h^2 - 2kdh - k^2 d^2 + 2(kd^2 + dh)\ell \\ &\quad - d^2 \ell^2 + n(h+kd) - \ell dn)], \\ A_3 &= \frac{1}{12cn} \sum_{\ell=k+1}^m h_{i,1+(\ell-1)d} [n^2 - 1 - 6(-h-kd+\ell d)(n+h+kd-\ell d)], \\ &= \frac{1}{12cn} \sum_{\ell=k+1}^m h_{i,1+(\ell-1)d} [n^2 - 1 - 6(-h^2 - 2kdh - k^2 d^2 + 2(kd^2 + dh)\ell \\ &\quad - d^2 \ell^2 - n(h+kd) + \ell dn)]. \end{aligned}$$

Besides,

$$\begin{aligned}
B_1 &= \sum_{\ell=2}^k h_{i,1+(\ell-1)d} = -\frac{d(k-1)}{n} - \frac{a}{2c} \frac{1}{(T_m(q)-1)} \left[(i-1) \left(U_{k-2}(q) + U_{m-k}(q) \right. \right. \\
&\quad \left. \left. - U_{m-1}(q) \right) - \frac{c}{a} \left(V_{k-1}(q) - V_{m-k}(q) + V_{m-1}(q) \right) - 1 \right], \\
B_2 &= \sum_{\ell=2}^k \ell h_{i,1+(\ell-1)d} = -\frac{d(k-1)(k+2)}{n} - \frac{a}{2c} \frac{1}{(T_m(q)-1)} \left[(i-1) \left((k+1)U_{k-2}(q) \right. \right. \\
&\quad \left. \left. + (k+1)U_{m-k}(q) - 2U_{m-2}(q) - 1 - \frac{1}{2(q-1)} (V_{k-1}(q) - V_{m-k}(q) + V_{m-1}(q)) \right) \right. \\
&\quad \left. - \frac{c}{a} \left(kV_{k-1}(q) - kV_{m-k}(q) + V_{m-1}(q) - U_{k-2}(q) - U_{m-k}(q) + U_{m-1}(q) \right) \right],
\end{aligned}$$

and

$$\begin{aligned}
K_1 &= \sum_{\ell=1}^m h_{i,1+(\ell-1)d} = 0, \\
K_2 &= \sum_{\ell=2}^k \ell h_{i,1+(\ell-1)d} = -\frac{1}{(T_m(q)-1)} \left[\left((i-1) \left(a - \frac{1}{d} - \frac{a}{2c} (m+2) \right) \frac{m+2}{2} \right) \right. \\
&\quad \left. \left(V_{m-1}(q) - 1 \right) + \left(\frac{a}{2c} (i-1)(m-1) - q + 1 \right) U_{m-1}(q) \right] - \frac{m+1}{2}, \\
K_3 &= \sum_{\ell=2}^k \ell^2 h_{i,1+(\ell-1)d} = -\frac{1}{(T_m(q)-1)} \left[- \left((i-1) \left(\frac{m+3}{d} + \frac{anm}{2cd} + \frac{an}{cd} \right) + \frac{m^2}{2} + m + 1 + \frac{2c}{ad} \right) \right. \\
&\quad \left. \left(V_{m-1}(q) - 1 \right) + \left(\frac{a}{2c} (i-1)(m^2 + 2m - 3)m - q \right) U_{m-1}(q) \right] - \frac{(m+1)(2m+1)}{2}.
\end{aligned}$$

Now adding $A_1 + A_2 + A_3 =$

$$\begin{aligned}
&= \frac{1}{12cn} (n^2 - 1) \sum_{\ell=1}^m h_{i,1+(\ell-1)d} - \frac{1}{2cn} h_{i,1} [-6(h + kd - d)(n - h - kd + d)] \\
&\quad - \frac{1}{2cn} \sum_{\ell=2}^m h_{i,1+(\ell-1)d} (-h^2 - 2kdh - k^2d^2 + 2(kd^2 + dh)\ell - d^2\ell^2) \\
&\quad - \frac{1}{2cn} \sum_{\ell=2}^k h_{i,1+(\ell-1)d} n(h + kd) - \ell dn + \frac{1}{2cn} \sum_{\ell=k+1}^m h_{i,1+(\ell-1)d} n(h + kd) - \ell dn \\
&\quad = -\frac{1}{2cn} \left[h_{i,1}(h + kd - d)(n - h - kd + d) + (-h^2 - 2kdh - k^2d^2)(-h_{i,1}) \right. \\
&\quad \quad + 2(kd^2 + dh)(K_2 - h_{i,1}) - d^2(K_3 - h_{i,1}) + n(h + kd)(2B_1 + h_{i,1}) \\
&\quad \quad \left. - dn(2B_2 - K_2 + h_{i,1}) \right] = -\frac{1}{2cn} \left[2nh_{i,1}(kd + h - d) + 2n(h + kd)B_1 \right. \\
&\quad \quad \left. - 2ndB_2 + (dn + 2dh + 2kd^2)K_2 - d^2K_3 \right] \\
&\quad - \frac{1}{2cn} \frac{1}{T_m(q) - 1} \left[n(h - i + 1)(V_{k-1}(q) - V_{m-k}(q)) \right. \\
&\quad \quad \left. - \left(\frac{an}{c}(i - 1)(h - d) - nd \right) (U_{k-2}(q) + U_{m-k}(q)) \right. \\
&\quad \quad \left. - \frac{2cd}{a} (V_{m-1}(q) - 1) - d^2U_{m-1}(q) \right] + dh + nh - nd - \frac{n^2}{6} + \frac{d^2}{6} \\
&\quad + (2kd - 3d + 2h - n)(i - 1) + kd(n - 2h) - d^2(k - 1)^2,
\end{aligned}$$

Finally, taking into account that $qU_{m-1}(q) - U_{m-2}(q) - 1 = T_m(q) - 1$, we obtain the desired result. \square

Proposition 2. *The group inverse of the Schur complement of \mathbf{L} is a block Toeplitz matrix given by*

$$\mathbf{M} = \begin{bmatrix} \mathbf{M}_1 & \mathbf{M}_2 & \dots & \mathbf{M}_m \\ \mathbf{M}_m & \mathbf{M}_1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \mathbf{M}_2 \\ \mathbf{M}_2 & \dots & \mathbf{M}_m & \mathbf{M}_1 \end{bmatrix}$$

where each submatrix \mathbf{M}_k has as elements

$$\begin{aligned}
(\mathbf{M}_k)_{i,h+1} &= \frac{1}{2cn} \left(-|i - 1 - k(d - 1) - h| \cdot (n - |i - 1 - k(d - 1) - h|) \right. \\
&\quad + \frac{1}{T_m(q) - 1} \left[n(h - i + 1)(V_{k-1}(q) - V_{m-k}(q)) \left(\frac{an}{c}(i - 1)(h - d) - nd \right) (U_{k-2}(q) + U_{m-k}(q)) \right. \\
&\quad \left. - \frac{2cd}{a} (V_{m-1}(q) - 1) - d^2U_{m-1}(q) \right] + dh + nh - nd + (2kd - 3d + 2h - n)(i - 1) + kd(n - 2h) \\
&\quad \left. - d^2(k - 1)^2 \right) + \frac{d^2 - 1}{12cn},
\end{aligned}$$

with $q = \frac{ad}{2c} + 1$, for any $k = 1, \dots, m$, $i = 1, \dots, d$ and $h = 0, \dots, d - 1$.

Lemma 6. *The product $\mathbf{Gs} = \mathbf{n} \otimes \mathbf{j}_m$ is a column vector where*

$$(\mathbf{n})_i = -\frac{a}{12cn} \left(n(6 + d) + 5m + 6mi(i - d - 2) \right), \text{ for any } i = 1, \dots, d.$$

Besides, it holds $\mathbf{Ms} = \mathbf{Gs}$.

Proof. For a given row $i = 1, \dots, d$, we have

$$\begin{aligned} (\mathbf{Gs})_i &= -a \sum_{k=1}^m g_{i,1+(k-1)d} = -ag_{i,1} - a \sum_{k=2}^m g_{i,1+(k-1)d} = -\frac{a}{12cn} (n^2 - 1)m \\ &\quad - \frac{a}{2cn} \left((i-1)(n-i+1) + \sum_{k=2}^m (1 + (k-1)d - i)(n-1 - (k-1)d + i) \right) \\ &= \frac{-a}{12cn} (6n + 5m + dn - 6mi(d+2-i)). \end{aligned}$$

Moreover, as \mathbf{G} is a block Toeplitz matrix, it holds $(\mathbf{Gs})_i = (\mathbf{Gs})_{i+(k-1)d}$, for any $k = 1, \dots, m$.

Besides, $\mathbf{Ms} = (\mathbf{G} - \mathbf{GFG})\mathbf{s} = \mathbf{Gs} - \mathbf{GFGs} = \mathbf{Gs}$, as it is straightforward to verify that $\mathbf{FGs} = (0, \dots, 0)^\top$.

□

5. ACKNOWLEDGEMENT

This work has been supported by the Spanish Research Council (Ministerio de Ciencia e Innovación) under the project MTM2014-60450-R.

REFERENCES

- [1] A. BEN-ISRAEL, T.N.E. GREVILLE: *Generalized inverses. Theory and applications*. 2nd Edition, Springer, New York, 2003.
- [2] M.C. GOLUMBIC, M. STERN, A. LEVY, G. MORGENSTERN: *Graph-Theoretic Concepts in Computer Science: 38th International Workshop, WG 2012, Jerusalem, Israel, June 26-28, 2012, Revised Selected Papers*. Springer, 2012 Edition.
- [3] E. BENDITO, A. CARMONA, A.M. ENCINAS, J.M. GESTO: *A Formula for the Kirchhoff Index*. Int. J. Quantum Chem., **108** (2008), 1200–1206.
- [4] E. BENDITO, A. CARMONA, A.M. ENCINAS, M. MITJANA: *Generalized inverses of symmetric M-matrices*. Linear Algebra Appl., **432** (2010), 2438–2454.
- [5] A. CARMONA, A.M. ENCINAS, M. MITJANA: *Discrete elliptic operators and their Green operators*. Linear Algebra Appl., **442** (2014), 115–134.
- [6] A. CARMONA, A.M. ENCINAS, S. GAGO, M.J. JIMÉNEZ, M. MITJANA: *The inverse of some circulant matrices*. Appl. Math. Comput., **270** (2015), 785–793.
- [7] A. CARMONA, A.M. ENCINAS, S. GAGO, M. MITJANA: *Green operators of networks with a new vertex*. Linear Algebra Appl., **491** (2016), 419–433.
- [8] Z. CINKIR: *Deletion and contraction identities for the resistance values and the Kirchhoff index*. Int. J. Quantum Chem., **111** (2011), 4030–4041.
- [9] C.M. DA FONSECA, J. PETRONILHO: *Explicit inverses of some tridiagonal matrices*. Linear Algebra Appl., **325** (2001), 7–21.
- [10] P.J. DAVIS: *Circulant Matrices*. John Wiley & sons, New York, (1979).
- [11] A.M. ENCINAS, M.J. JIMÉNEZ: *Floquet theory for second order linear homogeneous difference equations*. J. Dif. Eq. and App., (2016), DOI:10.1080/10236198.2015.1100609.
- [12] M.J. JIMÉNEZ: *Ecuaciones en diferencias lineales de segundo orden y su aplicación al Análisis matricial*. Phd. Thesis, (2016).
- [13] G. MEURANT: *A review on the inverse of symmetric tridiagonal and block tridiagonal matrices*. SIAM J. Matrix Anal. Appl., **13** (1992), 707–728.
- [14] W. WANG, D. YANG, Y. LUO: *The Laplacian polynomial and Kirchhoff index of graphs derived from regular graphs*. Discrete Appl. Math., **161** (2013), 3063–3071.

- [15] M.A. WOODBURY *Inverting modified matrices*, Statistical Research Group, Memo. Rep. no. 42, Princeton University, Princeton, N. J, 1950.